



Atmospheric Carbon Dioxide from Peat Wetland Ecosystems

A.E. Normand¹, B.L. Turner^{1,2}, A.N. Smith¹, B. Baiser¹, M.W. Clark¹, J.R. Long^{1,3}, S.P. Grover⁴, K.R. Reddy¹

1. University of Florida; 2. Smithsonian Tropical Research Institute, Panama;

3. National MagLab - AMRIS Facility; 4. La Trobe University, Australia



Funding Grants: G.S. Boebinger (NSF DMR-1644779); NSF Graduate Fellowship Anna Normand (GMO2432)


Peat soils release carbon into the atmosphere when land is drained for use or warmed by a changing climate. The potential for carbon dioxide (CO₂) gas formation by aerobic microbial conversion depends on the soil's organic carbon source. Soil content is determined by mean annual temperature, vegetation type, and land-use conditions.

In this research, soil samples were obtained from 125 global peatland ecosystems for analysis by solid-state nuclear magnetic resonance spectroscopy (ssNMR). A subset of 11 sites revealed that oxygen-alkyl chemistry (i.e., carbohydrates) are the strongest predictor of aerobic CO₂ production.


This research supports and extends previous studies of temperate and boreal peatlands that linked CO₂ production to polysaccharide or oxygen-alkyl carbon composition. This research also suggests that climate models can be improved by using oxygen-alkyl carbon content to predict risk of increased CO₂ production.

Facilities and instrumentation used: Peat samples were analyzed by magic angle spinning (MAS) ¹³C NMR spectroscopy at the MagLab's AMRIS Facility using a 3.2 mm E-free H/C/N probe built to a specialized MagLab probe design that protects the sample from chemical degradation.

Citation: Normand, A.E.; Turner, B.L.; Lamit, L.J.; Smith, A.N.; Baiser, B.; Clark, M.W.; Hazlett, C.; Kane, E.S.; Lilleskov, E.; Long, J.R.; Grover, S.P.; Reddy, K.R., *Organic matter chemistry drives carbon dioxide production of peatlands*, Geophysical Research Letters, **48** (18), e2021GL093392 (2021)
doi.org/10.1029/2021GL093392 – Links to Data [Set 1](#) and [Set 2](#)



Peatland locations and sites tested



Peat soils vary by mean annual temperature (−3 to 26°C); vegetation (mossy §, herbaceous †, or woody ‡), and land-use by humans (nonimpacted and long-term impacted*).

Spectral regions correspond to carbon functional groups: alkyl (0–45ppm), methoxyl (45–60ppm), oxygen-alkyl (60–110ppm), aromatic (110–160ppm), and carboxyl (160–220ppm).

