



Evidence of Striped Electronic Phases in a Structurally Modulated Superlattice

A. Devarakonda^{1,6}, A. Chen¹, D. Graf², M. Kriener³, A.J. Akey⁴, D.C. Bell⁴, T. Suzuki⁵, J.G. Checkelsky¹

1. MIT; 2. NHMFL; 3. RIKEN; 4. Harvard University; 5. Toho University; 6. Columbia University



Funding Grants: K. Amm (NSF DMR-2128556); J. Checkelsky (GBMF9070, DOE DE-SC0022028, ONR N00014-21-1-259); A. Devarakonda (Simons Foundation 855186); D.C. Bell (NSF DMR-1644779)

The electronic properties of materials (crystals) can be manipulated via the application of modulated electric & magnetic fields or via modulations engineered into the structure of the material itself. Similar to the beating sound produced by two tuning forks with slightly different frequencies, the spatial modulation produced between two crystal layers with slightly different (incommensurate) structures creates the conditions (stripes) for intriguing new electronic properties.

In layered SrTa_2S_5 such conditions are realized due to the difference in periodic structures of Sr_3TaS_5 and $H\text{-TaS}_2$ (Fig. 1a). These layers are formed as the material crystallizes from the molten base materials as the mixture is cooled in a furnace. The resulting modulation is $\sim 4.4\text{nm}$ in length which is quite long compared to the interatomic spacing in the crystal. Investigating the properties of SrTa_2S_5 revealed intriguing superconducting behavior which suggests that the first superconducting state forms at $T^* \sim 2.3\text{K}$ and is confined within the layers (Fig. 1b) due to the intralayer pairing of the electrons. Further cooling reveals a bulk superconducting state at $T_c = 1.49\text{K}$ with interlayer electron pairing (Fig. 1c).

To better understand why superconductivity behaves this way in SrTa_2S_5 measurements of the Fermi surface via quantum oscillations (QO) in resistance and magnetization were performed at the MagLab (Fig. 1d). The QO measurements revealed a complex superposition of frequencies from both the Fermi surface and oscillations resulting from the spatial modulation of the incommensurate interface between layers (Fig. 1e). The data paints a remarkable picture of electron mobility and coherent electron states in SrTa_2S_5 considering the low mobility and lack of QO in Sr_3TaS_5 & $H\text{-TaS}_2$ when measured individually.




Fig. 1: a) Crystal structure of SrTa_2S_5 , b) Illustration of *intralayer* electronic pairing, c) Illustration of *interlayer* electronic pairing, d) M vs. H traces to 31T showing quantum oscillations, e) Fast Fourier transform of the 5.3 deg. trace in d) showing component frequencies as well as the frequencies due to the modulated structure.

Facilities and instrumentation used: DC Field Facility, Cell 9 magnet (31T).

Citation: Devarakonda, A., Chen, A., Graf, D., Kriener, M., Akey, A.J., Bell, D.C., Suzuki, T., Checkelsky, J.G. *Evidence of striped electronic phases in a structurally modulated superlattice*, *Nature* (2024), doi.org/10.1038/s41586-024-07589-5