



A folding motif formed with an expanded genetic alphabet

Bang Wang^{1,2}, James R. Rocca², Zunyi Yang¹, Weihong Tan^{2,3}, Steven A. Benner¹, et al.
1. Foundation for Applied Molecular Evolution; 2. University of Florida; 3. Hunan University



Funding Grants: K. M. Amm (NSF DMR-2128556); S. A. Benner (NIH GM128186 and GM141391); W. Tan (National Natural Science Foundation of China T2188102)

This study explores how adding synthetic nucleotides to DNA not only increases its information density but also diversifies its three-dimensional structures. Specifically, the research introduces an additional nucleotide (dZ) to a DNA strand, forming a stable unimolecular structure called the folded Z-motif (fZ-motif). fZ potentially expands DNA technology applications in catalysis and information storage due to its compact and unique form.

Researchers used several analytical methods, including NMR spectroscopy, to characterize the fZ-motif. DNA strands containing the dZ nucleotide exhibit a stable and unique structure over a pH range of 8-9 with six reverse, skinny dZ:dZ⁻ base pairs. Their experiments highlight how the NMR capabilities of the MagLab can determine the unique structural properties of the fZ-motif and characterize the temperature and pH conditions over which it is stable.

The discovery of the fZ-motif advances the field of synthetic biology by demonstrating a new type of DNA folding enabled by an expanded genetic alphabet. This finding opens the door to potential applications in nanotechnology and molecular biology, including the development of new DNA-based sensors and nanomachines that respond to environmental changes.




Figure 1: Characterization of the fZ-motif formed by DNA containing dZ using ¹H NMR spectroscopy. A 2 mM solution of the ZZZ oligonucleotide was examined by NMR at 18.8T using aqueous buffers with varying pH and ¹H / ²H isotope ratios to characterize hydrogen exchange, a reporter on the stability of different hydrogen bonding patterns in DNA.

Facilities and instrumentation used: AMRIS Facility NMR Spectrometers at 600 MHz and 800 MHz (14.1 and 18.8T).

Citation: Wang, B.; Rocca, J.R.; Hoshika, S.; Chen, C.; Yang, Z.; Esmaeeli, R.; Wang, J.; Pan, X.; Lu, J.; Wang, K.; Cao, Y.; Tan, W.; Benner, S.A., *A folding motif formed with an expanded genetic alphabet*, *Nature Chemistry*, 1-10 (2024) doi.org/10.1038/s41557-024-01552-7